

A NOTE ON THE UNIQUENESS OF INVOLUTION IN LOCALLY C^* -ALGEBRAS

ALEXANDER A. KATZ

ABSTRACT. In the present note we show that the involution in locally C^* -algebras is uniquely determined.

1. INTRODUCTION

One of the important basic facts of the theory of C^* -**algebras** is that the unary operation of involution in a C^* -algebra is uniquely determined. This property was first observed in 1955 by Bohnenblust and Karlin in [2] (see as well [7] for a nice exposition).

The Hausdorff projective limits of projective families of Banach algebras as natural locally-convex generalizations of Banach algebras have been studied sporadically by many authors since 1952, when they were first introduced by Arens [1] and Michael [6]. The Hausdorff projective limits of projective families of C^* -algebras were first mentioned by Arens [1]. They have since been studied under various names by many authors. Development of the subject is reflected in the monograph of Fragoulopoulou [3]. We will follow Inoue [4] in the usage of the name **locally C^* -algebras** for these algebras.

The purpose of the present notes is to show that the unary operation of involution in locally C^* -algebras is uniquely determined.

2. PRELIMINARIES

First, we recall some basic notions on topological $*$ -algebras. A $*$ -algebra (or involutive algebra) is an algebra A over \mathbb{C} with an involution

$$* : A \rightarrow A,$$

such that

$$(a + \lambda b)^* = a^* + \bar{\lambda}b^*,$$

and

$$(ab)^* = b^*a^*,$$

for every $a, b \in A$ and $\lambda \in \mathbb{C}$.

A seminorm $\|\cdot\|$ on a $*$ -algebra A is a C^* -seminorm if it is submultiplicative, i.e.

$$\|ab\| \leq \|a\| \|b\|,$$

Date: January 2, 2012.

Key words and phrases. C^* -algebras, locally C^* -algebras, projective limit of projective family of C^* -algebras.

2010 AMS Subject Classification: Primary 46K05.

and satisfies the C^* -condition, i.e.

$$\|a^*a\| = \|a\|^2,$$

for every $a, b \in A$. Note that the C^* -condition alone implies that $\|\cdot\|$ is submultiplicative, and in particular

$$\|a^*\| = \|a\|,$$

for every $a \in A$ (cf. for example [3]).

When a seminorm $\|\cdot\|$ on a $*$ -algebra A is a C^* -norm, and A is complete in the topology generated by this norm, A is called a **C^* -algebra**. The following theorem is valid.

Theorem 1 (Bohnenblust and Karlin [2]). *The unary operation of involution in a C^* -algebra is uniquely determined.*

Proof. See for example [7] for details. \square

A topological $*$ -algebra is a $*$ -algebra A equipped with a topology making the operations (addition, multiplication, additive inverse, involution) jointly continuous. For a topological $*$ -algebra A , one puts $N(A)$ for the set of continuous C^* -seminorms on A . One can see that $N(A)$ is a directed set with respect to pointwise ordering, because

$$\max\{\|\cdot\|_\alpha, \|\cdot\|_\beta\} \in N(A)$$

for every $\|\cdot\|_\alpha, \|\cdot\|_\beta \in N(A)$, where $\alpha, \beta \in \Lambda$, with Λ being a certain directed set.

For a topological $*$ -algebra A , and $\|\cdot\|_\alpha \in N(A)$, $\alpha \in \Lambda$,

$$\ker \|\cdot\|_\alpha = \{a \in A : \|a\|_\alpha = 0\}$$

is a $*$ -ideal in A , and $\|\cdot\|_\alpha$ induces a C^* -norm (we as well denote it by $\|\cdot\|_\alpha$) on the quotient $A_\alpha = A / \ker \|\cdot\|_\alpha$, and A_α is automatically complete in the topology generated by the norm $\|\cdot\|_\alpha$, thus is a C^* -algebra (see [3] for details). Each pair $\|\cdot\|_\alpha, \|\cdot\|_\beta \in N(A)$, such that

$$\beta \succeq \alpha,$$

$\alpha, \beta \in \Lambda$, induces a natural (continuous) surjective $*$ -homomorphism

$$g_\alpha^\beta : A_\beta \rightarrow A_\alpha.$$

Let, again, Λ be a set of indices, directed by a relation (reflexive, transitive, antisymmetric) " \preceq ". Let

$$\{A_\alpha, \alpha \in \Lambda\}$$

be a family of C^* -algebras, and g_α^β be, for

$$\alpha \preceq \beta,$$

the continuous linear $*$ -mappings

$$g_\alpha^\beta : A_\beta \longrightarrow A_\alpha,$$

so that

$$g_\alpha^\alpha(x_\alpha) = x_\alpha,$$

for all $\alpha \in \Lambda$, and

$$g_\alpha^\beta \circ g_\beta^\gamma = g_\alpha^\gamma,$$

whenever

$$\alpha \preceq \beta \preceq \gamma.$$

Let Γ be the collections $\{g_\alpha^\beta\}$ of all such transformations. Let A be a $*$ -subalgebra of the direct product algebra

$$\prod_{\alpha \in \Lambda} A_\alpha,$$

so that for its elements

$$x_\alpha = g_\alpha^\beta(x_\beta),$$

for all

$$\alpha \preceq \beta,$$

where

$$x_\alpha \in A_\alpha,$$

and

$$x_\beta \in A_\beta.$$

Definition 1. The $*$ -algebra A constructed above is called a **Hausdorff projective limit** of the projective family

$$\{A_\alpha, \alpha \in \Lambda\},$$

relatively to the collection

$$\Gamma = \{g_\alpha^\beta : \alpha, \beta \in \Lambda : \alpha \preceq \beta\},$$

and is denoted by

$$\varprojlim A_\alpha,$$

and is called the *Arens-Michael decomposition* of A .

It is well known (see, for example [8]) that for each $x \in A$, and each pair $\alpha, \beta \in \Lambda$, such that $\alpha \preceq \beta$, there is a natural projection

$$\pi_\beta : A \longrightarrow A_\beta,$$

defined by

$$\pi_\alpha(x) = g_\alpha^\beta(\pi_\beta(x)),$$

and each projection π_α for all $\alpha \in \Lambda$ is continuous.

Definition 2. A topological $*$ -algebra A over \mathbb{C} is called a **locally C^* -algebra** if there exists a projective family of C^* -algebras

$$\{A_\alpha; g_\alpha^\beta; \alpha, \beta \in \Lambda\},$$

so that

$$A \cong \varprojlim A_\alpha,$$

i.e. A is topologically $*$ -isomorphic to a projective limit of a projective family of C^* -algebras, i.e. there exists its Arens-Michael decomposition of A composed entirely of C^* -algebras.

A topological $*$ -algebra A over \mathbb{C} is a locally C^* -algebra iff A is a complete Hausdorff topological $*$ -algebra in which topology is generated by a saturated separating family of C^* -seminorms (see [3] for details).

Example 1. Every C^* -algebra is a locally C^* -algebra.

Example 2. A closed $*$ -subalgebra of a locally C^* -algebra is a locally C^* -algebra.

Example 3. The product $\prod_{\alpha \in \Lambda} A_\alpha$ of C^* -algebras A_α , with the product topology, is a locally C^* -algebra.

Example 4. Let X be a compactly generated Hausdorff space (this means that a subset $Y \subset X$ is closed iff $Y \cap K$ is closed for every compact subset $K \subset X$). Then the algebra $C(X)$ of all continuous, not necessarily bounded complex-valued functions on X , with the topology of uniform convergence on compact subsets, is a locally C^* -algebra. It is well known that all metrizable spaces and all locally compact Hausdorff spaces are compactly generated (see [5] for details).

Let A be a locally C^* -algebra. Then an element $a \in A$ is called **bounded**, if

$$\|a\|_\infty = \{\sup \|a\|_\alpha, \alpha \in \Lambda : \|\cdot\|_\alpha \in N(A)\} < \infty.$$

The set of all bounded elements of A is denoted by $b(A)$.

It is well-known that for each locally C^* -algebra A , its set $b(A)$ of bounded elements of A is a locally C^* -subalgebra, which is a C^* -algebra in the norm $\|\cdot\|_\infty$, such that it is dense in A in its topology (see for example [3]).

3. THE UNIQUENESS OF INVOLUTION IN LOCALLY C^* -ALGEBRAS

Here we present the main theorem of the current notes.

Theorem 2. *The unary operation of involution in any locally C^* -algebra is unique, i.e., if $(A, *, \|\cdot\|_\alpha, \alpha \in \Lambda)$ and $(A, \#, \|\cdot\|_\alpha, \alpha \in \Lambda)$ are two locally C^* -algebras, means that each seminorm $\|\cdot\|_\alpha, \alpha \in \Lambda$, satisfies the C^* -property for both operations, " $*$ " and " $\#$ ", then*

$$* = \#$$

on A .

Proof. Let now A be a locally C^* -algebra, and let

$$A = \varprojlim A_\alpha,$$

$\alpha \in \Lambda$, be its Arens-Michael decomposition, built using the family of seminorms $\|\cdot\|_\alpha, \alpha \in \Lambda$, so that for each $\alpha \in \Lambda$,

$$(A_\alpha, *_\alpha, \|\cdot\|_\alpha)$$

and

$$(A_\alpha, \#_\alpha, \|\cdot\|_\alpha)$$

are C^* -algebras, where the unary operations " $*_\alpha$ " and " $\#_\alpha$ " on A_α are defined as follows:

$$\pi_\alpha(x^*) = (\pi_\alpha(x))^{*_\alpha},$$

and

$$\pi_\alpha(x^\#) = (\pi_\alpha(x))^{\#_\alpha},$$

for each $x \in A$ and $\alpha \in \Lambda$.

Let us now assume, to the contrary to the statement of the theorem, that there exists some $x \in A$, such that

$$x^* = y \neq z = x^\#.$$

Then there must exist $\alpha_0 \in \Lambda$, such that

$$\pi_{\alpha_0}(y) \neq \pi_{\alpha_0}(z).$$

In fact, if it is not the case, and

$$\pi_\alpha(y) = \pi_\alpha(z)$$

for each $\alpha \in \Lambda$, implies that

$$y = z,$$

which contradicts the assumption.

So, α_0 must be such that

$$\pi_{\alpha_0}(x^*) \neq \pi_{\alpha_0}(x^\#),$$

which means that for

$$\begin{aligned} \pi_{\alpha_0}(x) &= x_{\alpha_0} \in A_{\alpha_0}, \\ x_{\alpha_0}^{*\alpha} &\neq x_{\alpha_0}^{\#\alpha}, \end{aligned}$$

which contradicts Theorem 1. Found contradiction proves the theorem. \square

REFERENCES

- [1] **Arens, R.**, *A generalization of normed rings*. (English) Pacific J. Math., Vol. 2, 1952, pp. 455–471.
- [2] **Bohnenblust, H.F.; Karlin, S.**, *Geometrical properties of the unit sphere of Banach algebras*. (English) Ann. of Math. (2) No. 62, 1955, pp. 217–229.
- [3] **Fragoulopoulou, M.**, *Topological algebras with involution*. (English) North-Holland Mathematics Studies, Vol. 200, Elsevier Science B.V., Amsterdam, 2005, 495 pp.
- [4] **Inoue, A.**, *Locally C*-algebra*. (English), Mem. Fac. Sci. Kyushu Univ. Ser. A , Vol. 25, 1971, pp. 197–235.
- [5] **Kelley, J.L.**, *General topology*. (English) Reprint of the 1955 edition [Van Nostrand, Toronto, Ont.]. Graduate Texts in Mathematics, No. 27. Springer-Verlag, New York-Berlin, 1975, 298 pp.
- [6] **Michael, E.A.**, *Locally multiplicatively-convex topological algebras*. (English) Mem. Amer. Math. Soc., No. 11, 1952, 79 pp.
- [7] **Rickart, C.E.**, *General theory of Banach algebras*. (English) The University Series in Higher Mathematics D. van Nostrand Co., Inc., Princeton, N.J.-Toronto-London-New York, 1960, 394 pp.
- [8] **Trèves, F.**, *Topological vector spaces: Distributions and Kernels*. (English), New York-London: Academic Press., 1967, 565 pp.

DR. ALEXANDER A. KATZ, DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, ST. JOHN'S COLLEGE OF LIBERAL ARTS AND SCIENCES, ST. JOHN'S UNIVERSITY, 300 HOWARD AVENUE, DASILVA ACADEMIC CENTER 314, STATEN ISLAND, NY 10301, USA
E-mail address: `katza@stjohns.edu`